

Vermont Mathematics Talent Search, Solutions to Test 2, 2014-2015

Test and Solutions by Jean Ohlson and Evan Dummit

December 7, 2014

1. Three numbers form a geometric sequence. The arithmetic mean of the first two is -9, and the arithmetic mean of the first and third terms is -15. Find the smallest possible value of the first term.

Answer: -27 .

Solution: Let the three terms (in order) be a , ar , ar^2 . The given information states $\frac{a+ar}{2} = -9$ and $\frac{a+ar^2}{2} = -15$.

Rearranging gives $a(1+r) = -18$ and $a(1+r^2) = -30$. Solving the first equation for a yields $a = -\frac{18}{1+r}$, and plugging into the second equation yields $-\frac{18}{1+r}(1+r^2) = -30$, so $-18 - 18r^2 = -30r$.

Rearranging yields $18r^2 - 30r + 18 = 0$, so factoring yields $6(3r+1)(r-2) = 0$. Then $(r, a) = (2, -6)$ or $(-1/3, -27)$. The smallest possible value for a is $\boxed{-27}$.

2. A circular sector has the same perimeter and area as a rectangle. (The perimeter of the sector includes the two radii.) Prove that the radius of the sector equals one of the side lengths of the rectangle.

Solution 1: Let the rectangle have sides a and b , and the sector have radius r and arclength $2c$. Then we have $2a+2b = 2r+2c$ and $ab = rc$. This is equivalent to $a+b = r+c$ and $ab = rc$, so, if x is a variable, then we see $(x-a)(x-b) = (x-r)(x-c)$. Since a quadratic polynomial's roots are unique, we conclude either that $r = a$ or $r = b$.

Solution 2: Let the rectangle have sides a and b , and let the sector have radius r and angle θ . Then $2a+2b = 2r+r\theta$ and $ab = \frac{1}{2}r^2\theta$. Solving the first equation for θ yields $\theta = 2\frac{a+b-r}{r}$, and then plugging into the second equation yields $ab = \frac{1}{2}r^2 \cdot 2\frac{a+b-r}{r} = r(a+b-r)$. Rearranging and factoring yields $(r-a)(r-b) = 0$, so $r = a$ or $r = b$.

3. If Q is a convex quadrilateral whose four side lengths and two diagonal lengths all lie in the set $\{1, d\}$, where $d > 1$, find all possible values of d .

Answer: $d = \sqrt{2}, \sqrt{3}, \frac{1+\sqrt{5}}{2}, \frac{\sqrt{2}+\sqrt{6}}{2}$. (The last value can equivalently be written as $\sqrt{2+\sqrt{3}}$.)

Solution: There are several cases:

- All four sides of the quadrilateral have length 1, and the two diagonals have length d . In this case, Q is a square, and $d = \sqrt{2}$.
- All four sides of the quadrilateral and one diagonal have length 1, and the other diagonal has length d . In this case, Q is a rhombus obtained by gluing two equilateral triangles together along a side, so $d = \sqrt{3}$.
- Three sides of the quadrilateral have length 1, and one side has length d . In this case, it is not possible for either diagonal to have length 1, so they must both have length d as well. Then the quadrilateral is centrally symmetric, so it is a trapezoid $ABCD$ with base AD . If the diagonals intersect at E , we can calculate that $AE = DE = 1$, and so by similar triangles we observe that $d = \frac{1+\sqrt{5}}{2}$. (In fact, this trapezoid is what is obtained by removing one vertex from a regular pentagon.)

- (d) The quadrilateral's sides (in order) are $\{1, d, 1, d\}$. In this case, Q is a parallelogram. However, this cannot actually occur, because the long diagonal necessarily has length $> d$.
- (e) The quadrilateral's sides (in order) are $\{1, 1, d, d\}$. In this case, Q is a kite: say $ABCD$ where $AB = AC = 1$, and $BD = CD = d$, where the diagonals AD and BC intersect at E . If $BC = 1$, then $AE = \sqrt{3}/2$ and $DE = \sqrt{d^2 - 1}/4$, so since the sum must exceed 1, it would be the case that $\frac{\sqrt{3}}{2} + \sqrt{d^2 - \frac{1}{4}} = d$. However, the square of the left-hand side is $\frac{3}{4} + \sqrt{3}\sqrt{d^2 - \frac{1}{4}} + d^2 - \frac{1}{4} \geq d^2 + \frac{1}{2}$, so it cannot equal d . Thus $BC = d$, and then $AE = \sqrt{1 - d^2/4} + d\frac{\sqrt{3}}{2}$. The square of this sum must exceed 1, so it is equal to d . Then we obtain $\sqrt{1 - \frac{d^2}{4}} = d\left(1 - \frac{\sqrt{3}}{2}\right)$, so $d^2 = 2 + \sqrt{3} = \frac{4 + 2\sqrt{3}}{2}$.
- Taking the square root yields $d = \sqrt{2 + \sqrt{3}} = \frac{\sqrt{2} + \sqrt{6}}{2}$.
- (f) Three or four of the quadrilateral's sides have length d . In this case, if A is a vertex with an obtuse angle and B, C are adjacent vertices, then at least one of AB and AC has length d . Then BC (being opposite the obtuse angle A) would thus necessarily have length $> d$, which is impossible.

Remark: If the quadrilateral Q is not required to be convex, the two values $d = \sqrt{3}$ and $d = \frac{\sqrt{2} + \sqrt{6}}{2}$ can also occur in a non-convex quadrilateral, but there are no additional values of d obtained in these cases.

4. The remainder when dividing the polynomial $p(x)$ by $x^3 - 2x^2 - x + 2$ is $ax^2 + 6x + 12$. The remainder when dividing $p(x)$ by $x^3 + x^2 - 4x - 4$ is $3x^2 - bx - 12$. Find $a + b$.

Answer: -3 .

Solution: By hypothesis, there exist polynomials $s(x)$ and $t(x)$ such that

$$\begin{aligned} q(x) &= (x^3 - 2x^2 - x + 2)s(x) + (ax^2 + 6x + 12) \\ q(x) &= (x^3 + x^2 - 4x - 4)t(x) + (3x^2 - bx - 12). \end{aligned}$$

Setting $x = -1$ yields

$$\begin{aligned} q(-1) &= 0 \cdot s(-1) + (a + 6) \\ q(-1) &= 0 \cdot t(-1) + (b - 9) \end{aligned}$$

from which we deduce that $a + 6 = 3b - 9$, so that $b = a + 15$. Similarly, setting $x = 2$ yields

$$\begin{aligned} q(-2) &= 0 \cdot s(2) + (4a + 24) \\ q(-2) &= 0 \cdot t(2) + (-2b) \end{aligned}$$

from which we deduce that $4a + 24 = -2b = -2(a + 15)$, from which we obtain $a = -9$ and thus $b = 6$. Then $a + b = \boxed{-3}$.

Remark One such polynomial is $p(x) = -4x^3 - x^2 + 10x + 4$, whose remainder upon dividing by $x^3 + x^2 - 4x - 4$ is $-9x^2 + 6x + 12$ and whose remainder upon dividing by $x^3 - 2x^2 - x + 2$ is $3x^2 - 6x - 12$.

5. The "run-length" of a sequence of heads and tails is the length of the longest consecutive sequence of identical outcomes: thus, the run-length of the sequence HTTHHHHTH is 4, while the run-length of the sequence TTTHHTHTTT is 3. If a fair coin is flipped 12 times, find the probability that the run-length of the sequence of outcomes is 2.

Answer: $29/256$.

Solution: Clearly, there are exactly 2 sequences of run-length 1, as such a sequence must alternate heads and tails. Now we count sequences of run length at most 2. There are four types of such sequences: ones that end in a run of 1H, a run of 2H, a run of 1T, and a run of 2T.

Let a_n be the number of sequences ending in one H, b_n be the number of sequences ending in HH, c_n be the number of sequences ending in T, and d_n be the number of sequences ending in TT. (We implicitly assume that if there is a flip before the given sequence H, HH, T, or TT, then it is of the opposite kind.)

A sequence of length ending in H is obtained by appending an H to a sequence ending in a T. Thus, $a_n = c_{n-1} + d_{n-1}$.

A sequence of length ending in HH is obtained by appending an H to a sequence ending in one H. Thus, $b_n = a_{n-1}$.

A sequence of length ending in T is obtained by appending a T to a sequence ending in an H. Thus, $c_n = a_{n-1} + b_{n-1}$.

A sequence of length ending in TT is obtained by appending a T to a sequence ending in one T. Thus, $d_n = c_{n-1}$.

We can also check easily that $a_1 = c_1 = 1$ and $b_1 = d_1 = 0$. We then recursively calculate

n	1	2	3	4	5	6	7	8	9	10	11	12
a_n	1	1	2	3	5	8	13	21	34	55	89	144
b_n	0	1	1	2	3	5	8	13	21	34	55	89
c_n	1	1	2	3	5	8	13	21	34	55	89	144
d_n	0	1	1	2	3	5	8	13	21	34	55	89

and so the total number of such sequences is $2(144 + 89) = 466$. Then there are 464 sequences of run-length 2, so the probability is $\frac{464}{2^{12}} = \frac{29}{256}$.

Remark It is easy to show by induction that $a_n = c_n = F_n$ and $b_n = d_n = F_{n-1}$, where F_n is the n th Fibonacci number. (There are other counting arguments that show this more directly.) Then the total number of possible sequences of run-length 2 of length n is $2F_{n+1} - 2$.

6. Suppose that $1 < a < b < c < d < e$ are positive integers such that $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d} + \frac{1}{e} = 1$. If e is equal to 3 times a prime number, find the sum of all possible values for d .

Answer: 34. $[(a, b, c, d, e) = (2, 4, 10, 12, 15) \text{ or } (2, 3, 11, 22, 33)].$

Solution: Suppose $e = 3p$. Imagine rearranging the equation $\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d} + \frac{1}{e} = 1$ to collect all the terms with denominator divisible by p on one side, and the terms with denominator not divisible by p on the other side, and then summing both quantities. Since one side has a denominator not divisible by p , we see that the sum of all terms with denominator divisible by p must (when reduced to lowest terms) have denominator not divisible by p . In particular, since $e = 3p$ is the largest term, the only possibilities for the collection of terms of denominator divisible by p are $\frac{1}{p} + \frac{1}{3p} = \frac{4}{3p}$, $\frac{1}{2p} + \frac{1}{3p} = \frac{5}{6p}$, or $\frac{1}{p} + \frac{1}{2p} + \frac{1}{3p} = \frac{11}{6p}$. We conclude that the only possibilities are $p = 2$, $p = 5$, or $p = 11$, since p must divide the numerator of the sum.

- Observe that $p = 2$ does not work, because $e = 6$ would require $a = 2$, $b = 3$, $c = 4$, $d = 5$, which does not satisfy the equation.
- If $p = 5$, then two terms must be $\frac{1}{10} + \frac{1}{15} = \frac{1}{6}$. The other three terms must have $\frac{1}{f} + \frac{1}{g} + \frac{1}{h} = \frac{5}{6}$ with $f < g < h < 15$. If $f \geq 3$ then $\frac{1}{f} + \frac{1}{g} + \frac{1}{h} \leq \frac{1}{3} + \frac{1}{4} + \frac{1}{5} = \frac{47}{60} < \frac{5}{6}$, so we must have $f = 2$. Then $\frac{1}{g} + \frac{1}{h} = \frac{1}{3}$: since $g = 6$ is too large and $g = 5$ does not work, we see that $(g, h) = (4, 12)$ is the only possibility. This gives the unique solution $(a, b, c, d, e) = (2, 4, 10, 12, 15)$, with $d = 12$.
- If $p = 11$, then three terms are $\frac{1}{11} + \frac{1}{22} + \frac{1}{33} = \frac{1}{6}$. The other two terms satisfy $\frac{1}{f} + \frac{1}{g} = \frac{5}{6}$ and it is clear that the only solution is $(f, g) = (2, 3)$. This gives the solution $(a, b, c, d, e) = (2, 3, 11, 22, 33)$, with $d = 22$.

We conclude that the only solutions are $(2, 4, 10, 12, 15)$ and $(2, 3, 11, 22, 33)$, so the sum of all possible d is 34.